

# Oxygen Fugacity in the Laboratory and in Terrestrial Systems

E. Woermann<sup>1</sup>, G.C. Ulmer<sup>2</sup>, G. Eriksson<sup>3</sup> and S.K. Saxena<sup>4</sup>

<sup>1</sup>Inst.f.Kristallographie, RWTH, D-52056 Aachen, Germany

<sup>2</sup>Geology Dept., Temple University, Philadelphia, Pa.

<sup>3</sup>GTT-Technologies, D-52134 Herzogenrath, Germany,

<sup>4</sup>CeSMEC, University Park, Miami, Fl.

## ***Abstract***

Several different synthetic phase assemblages – so called oxygen buffers – were devised by Eugster, 1957 in order to adjust defined oxygen fugacities in laboratories for experimental petrology at given temperatures and pressures. The “oxygen buffers”– and the corresponding acronyms – soon became popular and are now applied universally also to express  $f_{O_2}$  levels in natural systems where, however, a correlation with the phase assemblages of the originally defined buffers is not evident. A typical example is presented by the numerous analyses of basalts with oxygen fugacities of (QFM) (see Ulmer *et al.*, 2007).

From the fact that the  $f_{O_2}$ -data of the majority of basalts worldwide are arranged in a narrow belt along QFM (see Basaltic Volcanism Study Project, 1981) it is concluded that their redox state must be buffered. What may the properties and the quality of the buffers involved be? It must be realized that

–an oxygen buffer is a phase assemblage containing a certain element in different oxidation states. The entire periodic system was scanned and elements capable of coexisting in various oxidation states were marked.

–any buffer assemblage may react with any other one until one phase from buffer assemblage is quantitatively consumed and the latter buffer is thus annihilated. Eventually thus only one complete buffer assemblage can survive and will thus be dominating.

From the conditions cited above it is concluded that only the reaction  $2 H_2O = 2 H_2 + O_2$  may provide the required buffer. By thermodynamic calculations the oxygen fugacity of dissociating pure water close to QFM is confirmed for temperatures above 800 K, kinetically above the closing temperature of a

reaction between basaltic magma and water, *i. e.* for conditions prevailing for the eruption of MORBs.

The correlation between the oxidation state of a rock and the activity of water during its formation may be extended to further systems. Thus the fact that analyses of Nakhrites were showing an oxidation state close to QFM indicates that during their formation on Mars water must have been involved – although at present neither water nor hydrous components in Mars minerals directly seem to support this conclusion.

#### REFERENCES:

- Eugster, H.P.,(1957), Heterogeneous reactions involving oxidation and reduction at high pressures and temperatures. *J.Chem.Phys.* **26**, 1760–1761
- Ulmer, G.C. and Woermann, E.(2007), Thermodynamic price tags for a wet mantle. *Goldschmidt 2007* (this volume).
- Basaltic Volcanism Study Project*, 1981